

Probability Distribution in the SABR Model of Stochastic Volatility

Patrick Hagan

Bloomberg LP
499 Park Avenue
New York, NY 10022

Andrew Lesniewski

Ellington Management Group
53 Forest Avenue
Old Greenwich, CT 06870

Diana Woodward

Societe Generale
1221 Avenue of the Americas
New York NY 10020

Draft of June 3, 2004

Abstract

We study the SABR model of stochastic volatility [8]. This model is essentially an extension of Dupire's local volatility model, in which a suitable volatility parameter is assumed to be stochastic. SABR model allows for a large variety of shapes of volatility smiles, and it performs remarkably well in the swaptions and caps / floors markets. We refine the results of [8] by constructing an accurate and efficient asymptotic form of the probability distribution of forwards. We also discuss the impact of boundary conditions at zero forward on the volatility smile. Our analysis is based on a WKB type expansion for the heat kernel of a perturbed Laplace-Beltrami operator on a suitable Riemannian manifold.

Contents

1	Introduction	2
2	SABR model	5
2.1	Underlying process	5
2.2	Green's function	6
2.3	Boundary conditions at $\mathbf{x} = \mathbf{0}$	8
2.4	Solving the initial value problem	10
3	Stochastic geometry of the state space	11
3.1	SABR plane	11
3.2	Brownian motion on the SABR plane	14
3.3	Laplace-Beltrami operator on the SABR plane	15
4	Asymptotic expansion	17
5	Volatility smile	19
5.1	Marginal transition probability	19
5.2	Implied volatility	20
5.3	Impact of boundary conditions at zero forward	21
5.4	Explicit models	21
A	Heat equation on the Poincare plane	21
A.1	Lower bound on the Laplace-Beltrami operator	22
A.2	Construction of the Green's function	22
B	Some asymptotic expansions	25
B.1	Asymptotics of the McKean kernel	25
B.2	Laplace's method	26
B.3	Normalizing the SABR distribution	29

1 Introduction

The SABR model [8] of stochastic volatility attempts to capture the dynamics of smile in the interest rate derivatives markets which are dominated by caps / floors and swaptions. It provides a parsimonious, accurate, intuitive, and easily implementable framework for pricing, risk management, and relative value in those markets. The model describes the dynamics of a single forward (swap or LIBOR) rate

with stochastic volatility. The dynamics of the model is characterized by a function $b(f)$ of the forward rate f which determines the general shape of the volatility skew, a parameter v which controls the level of the volatility of volatility, and a parameter ρ which governs the correlation between the changes in the underlying forward rate and its volatility. It is an extension of Black's model: choosing $v = 0$ and $b(f) = f$ reduces SABR to the lognormal Black model, while $v = 0$ and $b(f) = 1$ reduces it to the normal Black model.

The main reason why the SABR model has proven effective in the industrial setting is that, even though it is too complex to allow for a closed form solution, it has an accurate asymptotic solution. This solution, as well as its implications for pricing and risk management of interest derivatives, has been described in [8].

In this paper we refine the results presented in [8]. Our way of thinking has been strongly influenced by the asymptotic techniques which go by the names of the *geometric optics* or the *WKB method*, and, most importantly, by the classical results of Varadhan [17], [18] (see also [16], [12] for more recent presentations and refinements). These techniques allow one to relate the short time asymptotics of the fundamental solution (or the *Green's function*) of Kolmogorov's equation to the differential geometry of the state space. From the probabilistic point of view, the Green's function represents the transition probability of the diffusion, and it thus carries all the information about the process.

Specifically, let \mathcal{U} denote the state space of an n -dimensional diffusion process with no drift, and let $G_X(s, x)$, $x, X \in \mathcal{U}$, denote the Green's function. We also assume that the process is time homogeneous, meaning that the diffusion matrix is independent of s . Then, Varadhan's theorem states that

$$\lim_{s \rightarrow 0} s \log G_X(s, x) = -\frac{d(x, X)^2}{2}.$$

Here $d(x, X)$ is the geodesic distance on \mathcal{U} with respect to a Riemannian metric which is determined by the coefficients of the Kolmogorov equation. This gives us the leading order behavior of the Green's function. To extract usable asymptotic information about the transition probability, more accurate analysis is necessary, but the choice of the Riemannian structure on \mathcal{U} dictated by Varadhan's theorem turns out to be key. Indeed, that Riemannian geometry becomes an important book keeping tool in carrying out the calculations, rather than merely fancy language. Technically speaking, we are led to studying the asymptotic properties of the perturbed Laplace - Beltrami operator on a Riemannian manifold.

In order to explain the results of this paper we define a universal function $D(\zeta)$:

$$D(\zeta) = \log \frac{\sqrt{\zeta^2 - 2\rho\zeta + 1} + \zeta - \rho}{1 - \rho},$$

where ζ is the following combination of today's forward rate f , strike F , and a volatility parameter σ (which is calibrated so that the at the money options prices match the market prices):

$$\zeta = \frac{v}{\sigma} \int_F^f \frac{du}{b(u)}.$$

The function $D(\zeta)$ represents a certain metric whose precise meaning is explained in the body of the paper. The key object from the point of view of option pricing is the probability distribution of forwards $P_F(\tau, f)$. Our main result in this paper is the explicit asymptotic formula:

$$P_F(\tau, f) = \frac{\exp\left\{-D(\zeta)^2/2\tau v^2\right\}}{\sqrt{2\pi\tau}\sigma b(K)(\cosh D(\zeta) - \rho \sinh D(\zeta))^{3/2}} (1 + \dots).$$

In order not to burden the notation, we have written down the leading term only; the complete formula is stated in Section 5. To leading order, the probability distribution of forwards in the SABR model is Gaussian with the metric $D(\zeta)$ replacing the usual distance.

From this probability distribution, we can deduce explicit expressions for implied volatility. The normal volatility is given by:

$$\sigma_n = \frac{v(f-F)}{D(\zeta)} (1 + \dots),$$

while the lognormal volatility is given by:

$$\sigma_{ln} = \frac{v \log(f/F)}{D(\zeta)} (1 + \dots).$$

Precise formulas, including the subleading terms, are stated in Section 5.

We would like to mention that other stochastic volatility models have been extensively studied in the literature (notably among them the Heston model [11]). Useful presentations of these models are contained in [4] and [15]. We continue our approach to volatility modelling in [9].

A comment on our style of exposition in this paper. We chose to present the arguments in an informal manner. And while we believe that all the results of this paper could be stated and proved rigorously as theorems, little would be gained and clarity might easily get lost in the course of doing so.

The paper is organized as follows. In Section 2 we review the model and formulate the basic partial differential equation, the backward Kolmogorov equation. We also introduce the Green's and discuss various boundary conditions at zero. Section 3 is devoted to the description of the differential geometry underlying the SABR

model. We show that the stochastic dynamics defining the model can be viewed as a perturbation of the Brownian motion on a deformed Poincare plane. The elliptic operator in the Kolmogorov equation turns out to be a perturbed Laplace-Beltrami operator. This differential geometric setup is key to our asymptotic analysis of the model which is carried through in Section 4. In Section 5 we derive the explicit formulas for the probability distribution and implied volatility which we have discussed above. In Appendix A we review the derivation of the fundamental solution of the heat equation on the Poincare plane. This solution is the starting point of our perturbation expansion. Finally, Appendix B contains some useful asymptotic expansions.

2 SABR model

In this section we describe the SABR model of stochastic volatility [8]. It is a two factor model with the dynamics given by a system of two stochastic differential equations. The state variables of the model can be thought of as the forward price of an asset, and a volatility parameter. In order to derive explicit expressions for the associated probability distribution and the implied volatility, we study the Green's function of the backward Kolmogorov operator.

2.1 Underlying process

We consider a European option on a forward asset expiring T years from today. The forward asset that we have in mind can be for instance a forward LIBOR rate, a forward swap rate, or the forward yield on a bond. The dynamics of the forward in the SABR model is given by:

$$dF_t = \sigma_t b(F_t) dW_t, \quad (1)$$

$$d\sigma_t = v\sigma_t dZ_t. \quad (2)$$

Here F_t is the forward rate process, and W_t and Z_t are Brownian motions with

$$E[dW_t dZ_t] = \rho dt, \quad (3)$$

where the correlation ρ is assumed constant. Note that we assume that a suitable numeraire has been chosen so that F_t is a martingale. The process σ_t is the stochastic component of the volatility of F_t , and v is the volatility of σ_t (the “volga”) which is also assumed to be constant. The function $b(f)$ is defined for $f > 0$, and is assumed to be positive, monotone non-decreasing, and smooth. Two examples of b ,

which are particularly popular among financial practitioners, are functions of the form:

$$b(x) = x^\beta, \text{ where } 0 \leq \beta \leq 1 \quad (4)$$

(stochastic CEV model), or

$$b(x) = x + a, \text{ where } a > 0 \quad (5)$$

(stochastic shifted lognormal model).

The process σ_t is purely lognormal and thus $\sigma_t > 0$ almost surely. Since, depending on the choice of $b(x)$, the process F_t can take on positive only or positive and negative values with non-zero probability, we should carefully study the boundary behavior of the process (1)-(2), as F_t approaches 0. To this end, we extend the function $b(x)$ to all values of x by setting

$$b(-x) = -b(x), \text{ for } x < 0. \quad (6)$$

The so extended $b(x)$ is an odd function, $b(-x) = -b(x)$, for all values of x , and thus the process (1)-(2) is invariant under the reflection $F_t \rightarrow -F_t$. The state space of the extended process is thus the upper half plane.

Our analysis uses an asymptotic expansion in the parameter v^2T , and we thus require that v^2T be small. In practice, this is an excellent assumption for medium and longer dated options. Typical for shorter dated options are significant, discontinuous movements in implied volatility. The SABR model should presumably be extended to include such jump behavior of short dated options.

A special case of (1)-(2) which will play an important role in our analysis is the case of $b(x) = 1$, and $\rho = 0$. In this situation, the basic equations of motion have a particularly simple form:

$$dF_t = \sigma_t dW_t, \quad (7)$$

$$d\sigma_t = v\sigma_t dZ_t, \quad (8)$$

with $E[dW_t dZ_t] = 0$.

2.2 Green's function

Green's functions arise in finance as the prices of Arrow-Debreu securities. We consider the Arrow-Debreu security whose payoff at time T is given by Dirac's delta function $\delta(F_T - F, \sigma_T - \Sigma)$. The time $t < T$ price $G = G_{T,F,\Sigma}(t, f, \sigma)$ of this security is the solution to the following parabolic partial differential equation:

$$\frac{\partial G}{\partial t} + \frac{1}{2}\sigma^2 \left(b(f)^2 \frac{\partial^2 G}{\partial f^2} + 2v\rho b(f) \frac{\partial^2 G}{\partial f \partial \sigma} + v^2 \frac{\partial^2 G}{\partial \sigma^2} \right) = 0, \quad (9)$$

with the terminal condition:

$$G_{T,F,\Sigma}(t, f, \sigma) = \delta(f - F, \sigma - \Sigma), \text{ at } t = T. \quad (10)$$

This equation should also be supplemented by a boundary condition at infinity such that G is financially meaningful. Since the payoff takes place only if the forward has a predetermined value in a finite amount of time, the value of the Arrow-Debreu security has to tend to zero as F and Σ become large:

$$G_{T,F,\Sigma}(t, f, \sigma) \rightarrow 0, \quad \text{as } F, \Sigma \rightarrow \infty. \quad (11)$$

Thus $G_{T,F,\Sigma}(t, f, \sigma)$ is a Green's function for (9). Once we have constructed it, we can price any European option. For example, the price $C_{T,K}(t, f, \sigma)$ of a European call option struck at K and expiring at time T can be written in terms of $G_{T,F,\Sigma}(t, f, \sigma)$ as

$$C_{T,K}(t, f, \sigma) = \int (F - K)^+ G_{T,F,\Sigma}(t, f, \sigma) dF d\Sigma, \quad (12)$$

where, as usual, $(F - K)^+ = \max(F - K, 0)$, and where the integration extends over the upper half plane $\{(F, \Sigma) \in \mathbb{R}^2 : \Sigma > 0\}$.

Note that the process (1)-(2) is time homogeneous, and thus $G_{T,F,\Sigma}(t, f, \sigma)$ is a function of the time to expiry $\tau = T - t$ only. Denoting

$$G_{F,\Sigma}(\tau, f, \sigma) \equiv G_{T,F,\Sigma}(t, f, \sigma),$$

and

$$C_K(\tau, f, \sigma) \equiv C_{T,K}(t, f, \sigma),$$

we can reformulate (9)-(10) as the initial value problem:

$$\frac{\partial G}{\partial \tau} = \frac{1}{2} \sigma^2 \left(b(f)^2 \frac{\partial^2 G}{\partial f^2} + 2\nu \rho b(f) \frac{\partial^2 G}{\partial f \partial \sigma} + v^2 \frac{\partial^2 G}{\partial \sigma^2} \right), \quad (13)$$

and

$$G_{F,\Sigma}(\tau, f, \sigma) = \delta(f - F, \sigma - \Sigma), \text{ at } \tau = 0. \quad (14)$$

Introducing the marginal probability distribution

$$P_F(\tau, f, \sigma) = \int_0^\infty G_{F,\Sigma}(\tau, f, \sigma) d\Sigma, \quad (15)$$

we can express the call price (12) as

$$C_K(\tau, f, \sigma) = \int_{-\infty}^\infty (F - K)^+ P_F(\tau, f, \sigma) dF. \quad (16)$$

This formula has the familiar structure, and the focus of our analysis will be to derive a useful expression for $P_F(\tau, f)$.

We will solve (13)-(14) and (15) by means of asymptotic techniques. In order to set up the expansion, it is convenient to introduce the following variables:

$$s = \tau/T, \quad x = f, \quad X = F, \quad y = \sigma/v, \quad Y = \Sigma/v,$$

and the rescaled Green's function:

$$K_{X,Y}(s, x, y) = vTG_{X,vY}(Ts, x, vy).$$

In terms of these variables, the initial value problem (13)-(14) can be recast as:

$$\begin{aligned} \frac{\partial K}{\partial s} &= \frac{1}{2}\varepsilon y^2 \left(b(x)^2 \frac{\partial^2 K}{\partial x^2} + 2\rho b(x) \frac{\partial^2 K}{\partial x \partial y} + \frac{\partial^2 K}{\partial y^2} \right), \\ K(0, x, y) &= \delta(x - X, y - Y), \end{aligned} \quad (17)$$

where $K = K_{X,Y}$, and

$$\varepsilon = v^2 T. \quad (18)$$

It will be assumed that ε is small and it will serve as the parameter of our expansion. The heuristic picture behind this idea is that the volatility varies slower than the forward, and the rates of variability of f and σ/v are similar. The time T defines the time scale of the problem, and thus s is a natural dimensionless time variable. Expressed in terms of the new variables, our problem has a natural differential geometric content which is key to its solution.

Finally, let us write down the equations above for the special case of $b(x) = 1$, and $\rho = 0$:

$$\begin{aligned} \frac{\partial K}{\partial s} &= \frac{1}{2}\varepsilon y^2 \left(\frac{\partial^2 K}{\partial x^2} + \frac{\partial^2 K}{\partial y^2} \right), \\ K(0, x, y) &= \delta(x - X, y - Y). \end{aligned} \quad (19)$$

We will show later that this initial value problem has a closed form solution.

2.3 Boundary conditions at $x = 0$

The problem as we have formulated it so far is not complete. Since the value of the forward rate should be positive¹, we have to specify a boundary condition for the Green's function at $x = 0$. Three commonly used boundary conditions are [7]:

¹Recent history shows that this is not always necessarily the case, but we regard such occurrences as anomalous.

- *Dirichlet (or absorbing) boundary condition.* We assume that the Green's function, denoted by $K_{X,Y}^D(s, x, y)$, vanishes at $x = 0$,

$$K_{X,Y}^D(s, 0, y) = 0. \quad (20)$$

- *Neumann (or reflecting) boundary condition.* We assume that the derivative of the Green's function at $x = 0$, normal to the boundary (and pointing outward), vanishes. Let $K_{X,Y}^N(s, x, y)$ denote this Green's function; then

$$\frac{\partial}{\partial x} K_{X,Y}^N(s, 0, y) = 0. \quad (21)$$

- *Robin (or mixed) boundary condition.* The Green's function, which we shall denote by $K_{X,Y}^R(s, x, y)$, satisfies the following condition. Given $\eta > 0$,

$$\left(-\frac{\partial}{\partial x} + \eta \right) K_{X,Y}^R(s, 0, y) = 0. \quad (22)$$

In this paper we will be concerned with the Dirichlet and Neumann boundary conditions only. Our task is tremendously simplified by the fact that the differential operator in (17) is invariant under the reflection $x \rightarrow -x$ of the upper half plane. This allows one to construct the desired Green's functions by means of the method of images. Let $K_{X,Y}(s, x, y)$ denote now the solution to (17) which ignores any boundary condition at $x = 0^2$. Then, one verifies readily that

$$K_{X,Y}^D(s, x, y) = K_{X,Y}(s, x, y) - K_{X,Y}(s, -x, y), \quad (23)$$

and

$$K_{X,Y}^N(s, x, y) = K_{X,Y}(s, x, y) + K_{X,Y}(s, -x, y) \quad (24)$$

are the solutions to the Dirichlet and Neumann problem, respectively.

Observe that the Green's functions corresponding to these different boundary conditions obey the following *conditioning inequalities*:

$$K^D \leq K \leq K^N. \quad (25)$$

Since the Dirichlet boundary condition corresponds to the stochastic process being killed at the boundary, the total mass of the Green's function is less than one:

$$\int K_{X,Y}^D(s, x, y) dx dy < 1. \quad (26)$$

²This is sometimes referred to as the Green's function with a *free boundary condition*.

The remaining probability is a Dirac's delta function at $x = 0$. On the other hand, for the free and Neumann boundary conditions,

$$\int K_{X,Y}(s, x, y) dx dy = \int K_{X,Y}^N(s, x, y) dx dy = 1, \quad (27)$$

and so they are *bona fide* probability distributions.

2.4 Solving the initial value problem

It is easy to write down a formal solution to the initial value problem (17). Let L denote the partial differential operator

$$L = \frac{1}{2}y^2 \left(b(x)^2 \frac{\partial^2}{\partial x^2} + 2\rho b(x) \frac{\partial^2}{\partial x \partial y} + \frac{\partial^2}{\partial y^2} \right) \quad (28)$$

supplemented by a suitable boundary condition at $x = 0$. Consider the one-parameter semigroup of operators

$$U(s) = \exp(s\varepsilon L). \quad (29)$$

Then U solves the following initial value problem:

$$\begin{aligned} \frac{\partial U}{\partial s} &= \varepsilon L U, \\ U(0) &= I, \end{aligned}$$

and thus the Green's function $K_{X,Y}(s, x, y)$ is the integral kernel of $U(s)$:

$$K_{X,Y}(s, x, y) = U(s)(x, y; X, Y). \quad (30)$$

In order to solve the problem (17) it is thus sufficient to construct the semigroup $U(s)$ and find its integral kernel. Keeping in mind that our goal is to find an explicit formula for $K_{X,Y}(s, x, y)$, the strategy will be to represent L as the sum

$$L = L_0 + V, \quad (31)$$

where L_0 is a second order differential operator with the property that

$$U_0(s) = \exp(s\varepsilon L_0) \quad (32)$$

can be represented in closed form. The operator V turns out to be a differential operator of first order, and we will treat it as a small perturbation of the operator L_0 . The semigroup $U(s)$ can now be expressed in terms of $U_0(s)$ and V as

$$U(s) = Q(s)U_0(s). \quad (33)$$

Here, the operator $Q(s)$ is given by the well known regular perturbation expansion:

$$Q(s) = I + \sum_{1 \leq n < \infty} \int_{0 \leq s_1 \leq \dots \leq s_n \leq s\varepsilon} e^{s_1 \text{ad}_{L_0}}(V) \dots e^{s_n \text{ad}_{L_0}}(V) ds_1 \dots ds_n, \quad (34)$$

where ad_{L_0} is the commutator with L_0 :

$$\text{ad}_{L_0}(V) = L_0 V - V L_0. \quad (35)$$

We will use the first few terms in the expansion above in order to construct an accurate approximation to the Green's function $K_{X,Y}(s, x, y)$:

$$Q(s) = I + s\varepsilon V + \frac{1}{2} (s\varepsilon)^2 (\text{ad}_{L_0}(V) + V^2) + O((s\varepsilon)^3). \quad (36)$$

We shall disregard the convergence issues associated with this series, and use it solely as a tool to generate an asymptotic expansion.

3 Stochastic geometry of the state space

In solving our model we find that the SABR model in the special case $\rho = 0$, $b(x) = 1$ is Brownian motion on the Poincare plane. Generally, when $\rho \neq 0$, or $b(x) \neq 1$, the model amounts to Brownian motion on a two dimensional manifold, the *SABR plane*, perturbed by a drift term. In this section we summarize a number of basic facts about the differential geometry of the state space of the SABR model. The fundamental geometric structure is that of the Poincare plane. We will show that the state space of the SABR model can be viewed as a suitable deformation of the Poincare geometry.

3.1 SABR plane

We begin by reviewing the Poincare geometry of the upper half plane which will serve as the standard state space of our model. For a full (and very readable) account of the theory the reader is referred to e.g. [1].

The *Poincare plane* (also known as the hyperbolic or Lobachevski plane) is the upper half plane $\mathbb{H}^2 = \{(x, y) : y > 0\}$ equipped with the Poincare line element

$$ds^2 = \frac{dx^2 + dy^2}{y^2}. \quad (37)$$

This line element comes from the metric tensor given by

$$h = \begin{pmatrix} \frac{1}{y^2} & 0 \\ 0 & \frac{1}{y^2} \end{pmatrix}. \quad (38)$$

The Poincare plane admits a large group of symmetries. We introduce complex coordinates on \mathbb{H}^2 , $z = x + iy$ (the defining condition then reads $\text{Im}z > 0$), and consider a Moebius transformation

$$z' = \frac{az + b}{cz + d}, \quad (39)$$

where a, b, c, d are real numbers with $ad - bc = 1$. We verify easily the following two facts.

- Transformation (39) is a biholomorphic map of \mathbb{H}^2 onto itself.
- The Poincare metric is invariant under (39).

As a consequence, the Lie group

$$SL(2, \mathbb{R}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{R}, ad - bc = 1 \right\} \quad (40)$$

acts holomorphically and isometrically on \mathbb{H}^2 . This symmetry group plays very much the same role in the hyperbolic geometry as the Euclidean group in the usual Euclidean geometry of the plane \mathbb{R}^2 .

Let $d(z, Z)$ denotes the geodesic distance between two points $z, Z \in \mathbb{H}^2$, $z = x + iy$, $Z = X + iY$, i.e. the length of the shortest path connecting z and Z . There is an explicit expression for $d(z, Z)$:

$$\cosh d(z, Z) = 1 + \frac{|z - Z|^2}{2yY}, \quad (41)$$

where $|z - Z|$ denotes the Euclidean distance between z and Z . In particular, if $x = X$, then $d(z, Z) = |\log(y/Y)|$.

Since $\det(h) = y^{-4}$, the invariant volume element on \mathbb{H}^2 is given by

$$\begin{aligned} d\mu_h(z) &= \sqrt{\det(h)} \, dx dy \\ &= \frac{dx dy}{y^2}. \end{aligned} \quad (42)$$

The state space associated with the general SABR model has a somewhat more complicated geometry. Let \mathbb{M}^2 denote the upper half plane $\{(x, y) : y > 0\}$, equipped with the following metric g :

$$g = \frac{1}{\sqrt{1 - \rho^2}} \begin{pmatrix} \frac{1}{y^2 b(x)^2} & -\frac{\rho}{y^2 b(x)} \\ -\frac{\rho}{y^2 b(x)} & \frac{1}{y^2} \end{pmatrix}. \quad (43)$$

This metric is a generalization of the Poincare metric: the case of $\rho = 0$ and $b(x) = 1$ reduces to the Poincare metric. In fact, the metric g is the pullback of the Poincare metric under a suitable diffeomorphism. To see this, we choose $p > 0$, and define a map $\phi_p : \mathbb{M}^2 \rightarrow \mathbb{H}^2$ by

$$\phi_p(z) = \left(\frac{1}{\sqrt{1-\rho^2}} \left(\int_p^x \frac{du}{b(u)} - \rho y \right), y \right), \quad (44)$$

where $z = (x, y)$. The Jacobian $\nabla\phi_p$ of ϕ_p is

$$\nabla\phi_p(z) = \begin{pmatrix} \frac{1}{\sqrt{1-\rho^2} b(x)} & -\frac{\rho}{\sqrt{1-\rho^2}} \\ 0 & 1 \end{pmatrix}, \quad (45)$$

and so $\phi_p^*h = g$, where ϕ_p^* denotes the pullback of ϕ_p . The manifold \mathbb{M}^2 is thus isometrically diffeomorphic with the Poincare plane. A consequence of this fact is that we have an explicit formula for the geodesic distance $\delta(z, Z)$ on \mathbb{M}^2 :

$$\begin{aligned} \cosh \delta(z, Z) &= \cosh d(\phi_p(z), \phi_p(Z)) \\ &= 1 + \frac{\left(\int_X^x \frac{du}{b(u)} \right)^2 - 2\rho(y-Y) \int_X^x \frac{du}{b(u)} + (y-Y)^2}{2(1-\rho^2)yY}, \end{aligned} \quad (46)$$

where $z = (x, y)$ and $Z = (X, Y)$ are two points on \mathbb{M}^2 . In the following, we will suppress the subscript p , as the choice of p will be clear from the context. Since $\det(g) = y^{-4}b(x)^{-2}$, the invariant volume element on \mathbb{M}^2 is given by

$$\begin{aligned} d\mu_h(z) &= \sqrt{\det(g)} dx dy \\ &= \frac{dx dy}{b(x)y^2}. \end{aligned} \quad (47)$$

It will be convenient to use invariant notation. Let $z^1 = x$, $z^2 = y$, and let $\partial_\mu = \partial/\partial z^\mu$, $\mu = 1, 2$, denote the corresponding partial derivatives. We denote the components of g^{-1} by $g^{\mu\nu}$, and use g^{-1} and g to raise and lower the indices: $z_\mu = g_{\mu\nu}z^\nu$, $\partial^\mu = g^{\mu\nu}\partial_\nu = \partial/\partial z_\mu$, where we sum over the repeated indices. Explicitly,

$$\begin{aligned} \partial^1 &= y^2 \left(b(x)^2 \partial_1 + \rho b(x) \partial_2 \right), \\ \partial^2 &= y^2 \left(\rho b(x) \partial_1 + \partial_2 \right). \end{aligned}$$

Consequently, the initial value problem (17) can be written in the following geometric form:

$$\begin{aligned}\frac{\partial}{\partial s} K_Z(s, z) &= \frac{1}{2} \varepsilon \partial^\mu \partial_\mu K_Z(s, z), \\ K_Z(0, z) &= \delta(z - Z).\end{aligned}\tag{48}$$

3.2 Brownian motion on the SABR plane

It is no coincidence that the SABR model leads to the Poincare geometry. Recall [12] that the Brownian motion on the Poincare plane is described by the following system of stochastic differential equations:

$$dX_t = Y_t dW_t, \tag{49}$$

$$dY_t = Y_t dZ_t, \tag{50}$$

with the two Wiener processes W_t and Z_t satisfying

$$E[dW_t dZ_t] = 0. \tag{51}$$

Comparing this with the special case of the SABR model (7)-(8), we see that (7)-(8) reduces to (49)-(50) once we have made the following identifications:

$$X_t = F_{v^2 t}, \tag{52}$$

$$Y_t = \frac{1}{v} \sigma_{v^2 t}, \tag{53}$$

and used the scaling properties of a Wiener process:

$$dW_{v^2 t} = v dW_t,$$

$$dZ_{v^2 t} = v dZ_t.$$

Note that the system (49)-(50) can easily be solved in closed form: its solution is given by

$$X_t = X_0 + Y_0 \int_0^t \exp\left(Z_s - \frac{s^2}{2}\right) dW_s, \tag{54}$$

$$Y_t = Y_0 \exp\left(Z_t - \frac{t^2}{2}\right). \tag{55}$$

Let us now compare the SABR dynamics with that of the diffusion on the SABR plane. In order to find the dynamics of Brownian motion on the SABR

plane we use the fact that there is a mapping (namely, (44)) of \mathbb{M}^2 into \mathbb{H}^2 . Using this mapping and Ito's lemma yields the following system

$$dX_t = \frac{1}{2} Y_t^2 b(X_t) b'(X_t) dt + Y_t b(X_t) dW_t, \quad (56)$$

$$dY_t = Y_t dZ_t, \quad (57)$$

with the two Wiener processes W_t and Z_t satisfying

$$E[dW_t dZ_t] = \rho dt. \quad (58)$$

Note that this is not exactly the SABR model dynamics. Indeed, one can regard the SABR model as the perturbation of the Brownian motion on the SABR plane by the drift term $-\frac{1}{2} Y_t^2 b(X_t) b'(X_t) dt$.

As in the case of the Poincare plane, it is possible to represent the solution to the system (56)-(57) explicitly:

$$\int_{X_0}^{X_t} \frac{du}{b(u)} = Y_0 \int_0^t \exp\left(Z_s - \frac{s^2}{2}\right) dW_s, \quad (59)$$

$$Y_t = Y_0 \exp\left(Z_t - \frac{t^2}{2}\right). \quad (60)$$

Parenthetically, we note that, within Stratonovich's calculus, (56)-(57) can be written as

$$dX_t = Y_t b(X_t) \circ dW_t,$$

$$dY_t = Y_t \circ dZ_t.$$

Therefore, the stochastic differential equations of the SABR model, if interpreted according to Stratonovich, describe the dynamics of Brownian motion on the SABR plane.

3.3 Laplace-Beltrami operator on the SABR plane

Recall that the Laplace-Beltrami operator Δ_g on a Riemannian manifold \mathcal{M} with metric tensor g is defined by

$$\Delta_g f = \frac{1}{\sqrt{\det g}} \frac{\partial}{\partial x^\mu} \left(\sqrt{\det g} g^{\mu\nu} \frac{\partial f}{\partial x^\nu} \right), \quad (61)$$

where f is a smooth function on \mathcal{M} . It is a natural generalization of the familiar Laplace operator to spaces with non-Euclidean geometry. Its importance for probability theory comes from the fact that it serves as the infinitesimal generator of Brownian motion on such spaces (see e.g. [5], [6], [12]).

In the case of the Poincare plane, the Laplace-Beltrami operator has the form:

$$\Delta_h = y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right). \quad (62)$$

As anticipated by our discussion in Section 3.2, this operator is closely related to the operator L in the special case of $b(x) = 1$, $\rho = 0$. In fact, in this case,

$$L = \frac{1}{2} \Delta_h, \quad (63)$$

and thus the problem (19) turns out to be the initial value problem the heat equation on \mathbb{H}^2 :

$$\begin{aligned} \frac{\partial K_Z}{\partial s} &= \frac{1}{2} \varepsilon \Delta_h K_Z, \\ K_Z(0, z) &= \delta(z - Z) \end{aligned} \quad (64)$$

The key fact is that the Green's function for this equation can be represented in closed form,

$$K_Z^h(s, z) = \frac{e^{-s\varepsilon/8} \sqrt{2}}{(2\pi s\varepsilon)^{3/2} Y^2} \int_{d(z, Z)}^{\infty} \frac{u e^{-u^2/2s\varepsilon}}{\sqrt{\cosh u - \cosh d(z, Z)}} du. \quad (65)$$

This formula was originally derived by McKean [14] (see also [12] and references therein). We have added the superscript h to indicate that this Green's function is associated with the Poincare metric. In Appendix A we outline an elementary derivation of this fact.

Let us now extend the discussion above to the general case. We note first that, except for the case of $b(x) = 1$, the operator $\partial^\mu \partial_\mu$ does not coincide with the Laplace-Beltrami operator Δ_g on \mathbb{M}^2 associated with the metric (43). It is, however, easy to verify that

$$\begin{aligned} \partial^\mu \partial_\mu f &= \Delta_g f - \frac{1}{\sqrt{\det g}} \frac{\partial}{\partial x^\nu} \left(\sqrt{\det g} g^{\mu\nu} \right) \frac{\partial f}{\partial x^\mu} \\ &= \Delta_g f - \frac{1}{\sqrt{1 - \rho^2}} y^2 b b' \frac{\partial f}{\partial x}, \end{aligned}$$

and thus

$$\begin{aligned} L &= \frac{1}{2} \Delta_g - \frac{1}{2\sqrt{1 - \rho^2}} y^2 b b' \frac{\partial}{\partial x} \\ &= L_0 + V, \end{aligned}$$

where L_0 is essentially the Laplace-Beltrami operator:

$$L_0 = \frac{1}{2} \Delta_g, \quad (66)$$

and $V(x)$ is lower order:

$$V = -\frac{1}{2\sqrt{1-\rho^2}} y^2 b(x) b'(x) \frac{\partial}{\partial x}. \quad (67)$$

Let us first focus on the Laplace-Beltrami operator Δ_g . The key property of the Laplace-Beltrami operator is its invariance under a diffeomorphism. In particular, this implies that

$$\Delta_g = \phi^{-1} \circ \Delta_h \circ \phi, \quad (68)$$

and, hence, the heat equation

$$\frac{\partial K}{\partial s} = \frac{1}{2} \varepsilon \Delta_g K.$$

on \mathbb{M}^2 can be solved in closed form! The Green's function $K_Z^g(s, z)$ of this equation is related to (65) by

$$K_Z^g(s, z) = \det(\nabla \phi(Z)) K_{\phi(Z)}^h(s, \phi(z)). \quad (69)$$

Explicitly,

$$K_Z^g(s, z) = \frac{e^{-s\varepsilon/8} \sqrt{2}}{(2\pi s\varepsilon)^{3/2} \sqrt{1-\rho^2} Y^2 b(X)} \int_\delta^\infty \frac{u e^{-u^2/2s\varepsilon}}{\sqrt{\cosh u - \cosh \delta}} du, \quad (70)$$

where $\delta = \delta(z, Z)$ is the geodesic distance (46) on \mathbb{M}^2 . This is the explicit representation of the integral kernel of the operator $U_0(s)$.

4 Asymptotic expansion

In principle, we have now completed our task of solving the initial value problem (19). Indeed, its solution is given by

$$K_Z(s, z) = Q(s) K_Z^g(s, z), \quad (71)$$

where $Q(s)$ is the perturbation expansion given by (34). In order to produce clear results that can readily be used in practice we perform now a perturbation expansion on the expression above. Our method allows one to calculate the Green's function of the model to the desired order of accuracy.

Let us start with the Green's function $K_Z^h(s, z)$ which is defined on the Poincare plane. In Appendix B we derived an asymptotic expansion (104) for the heat kernel on the Poincare plane. After rescaling as in (93), we arrive at

$$K_Z^h(s, z) = \frac{1}{2\pi\lambda Y^2} \exp\left(-\frac{d^2}{2\lambda}\right) \times \sqrt{\frac{d}{\sinh d}} \left(1 - \frac{1}{8} \left(\frac{d \coth d - 1}{d^2} + 1\right) \lambda + O(\lambda^2)\right),$$

where we have introduced a new variable,

$$\lambda = s\varepsilon. \quad (72)$$

We can now extend the expression to the general Green's function $K_Z^g(s, z)$. Using (69) or (70) we find that $K_Z^g(s, z)$ has the following asymptotic expansion:

$$K_Z^g(s, z) = \frac{1}{2\pi\lambda\sqrt{1-\rho^2}Y^2b(X)} \exp\left(-\frac{\delta^2}{2\lambda}\right) \times \sqrt{\frac{\delta}{\sinh \delta}} \left(1 - \frac{1}{8} \left(\frac{\delta \coth \delta - 1}{\delta^2} + 1\right) \lambda + O(\lambda^2)\right).$$

To complete the calculation in the case of general $b(x)$ we need to take into account the contribution to the Green's function coming from perturbation V defined in (67). Let us define the function:

$$q(z, Z) = \sinh \delta(z, Z) V \delta(z, Z) = -\frac{yb'(x)}{2(1-\rho^2)^{3/2}Y} \left(\int_X^x \frac{du}{b(u)} - \rho(y-Y)\right). \quad (73)$$

From (104) and (105),

$$K_Z(s, z) = (I + \lambda V) K_Z^g(s, z) = \frac{1}{\sqrt{1-\rho^2}Y^2b(X)} \left(K_Z(s, z) + \lambda \frac{q}{\sinh \delta} \frac{\partial}{\partial \delta} K_Z(s, z)\right), \quad (74)$$

which yields the following asymptotic formula for the Green's function:

$$K_Z(s, z) = \frac{1}{2\pi\lambda\sqrt{1-\rho^2}Y^2b(X)} \exp\left(-\frac{\delta^2}{2\lambda}\right) \sqrt{\frac{\delta}{\sinh \delta}} \times \left(1 - \frac{\delta}{\sinh \delta} q - \left(\frac{1}{8} + \frac{\delta \coth \delta - 1}{8\delta^2} - \frac{\delta \coth \delta - 1}{2\delta \sinh \delta} q\right) \lambda + O(\lambda^2)\right). \quad (75)$$

In a way, this is the central result of this paper. It gives us a precise asymptotic behavior of the Green's function of the SABR model, as $\lambda \rightarrow 0$.

5 Volatility smile

We are now ready to complete our analysis. Given the explicit form of the approximate Green's function, we can calculate (via another asymptotic expansion) the marginal probability distribution. Comparing the result with the normal probability distribution allows us to find the implied normal and lognormal volatilities, as functions of the model parameters. We conclude this section by deriving explicit formulas for the case of the CEV model $b(x) = x^\beta$ and the shifted lognormal model $b(x) = x + a$.

5.1 Marginal transition probability

We first integrate the asymptotic joint density over the terminal volatility variable Y to find the marginal density for the forward x :

$$\begin{aligned} P_X(s, x, y) &= \int_0^\infty K_Z(s, z) dY \\ &= \frac{1}{2\pi\lambda\sqrt{1-\rho^2}b(X)} \int_0^\infty \frac{dY}{Y^2} e^{-\delta^2/2\lambda} \sqrt{\frac{\delta}{\sinh \delta}} \times \\ &\quad \left(1 - \frac{\delta}{\sinh \delta} q - \left(\frac{1}{8} + \frac{\delta \coth \delta - 1}{8\delta^2} - \frac{\delta \coth \delta - 1}{2\delta \sinh \delta} q \right) \lambda \right). \end{aligned} \quad (76)$$

Here the metric $\delta(z, Z)$ is defined implicitly by (46). We evaluate this integral asymptotically by using Laplace's method (steepest descent). This analysis is carried out in Appendix B.2. The key step is to analyze the argument Y of the exponent

$$\phi(Y) = \frac{1}{2} \delta(z, Z)^2, \quad (77)$$

in order to find the point Y_0 where this function is at a minimum. Let us introduce the notation:

$$\zeta = \frac{1}{y} \int_X^x \frac{du}{b(u)}.$$

Since $yb(u)$ is basically the rescaled volatility at forward u , $1/\zeta$ represents the average volatility between today's forward x and at option's strike X . In other words, ζ represents how "easy" it is to reach the strike X . Some algebra shows that the minimum of (77) occurs at $Y_0 = Y_0(\zeta, y)$, where

$$Y_0 = y\sqrt{\zeta^2 - 2\rho\zeta + 1}. \quad (78)$$

Also, let $D(\zeta)$ denote the value of $\delta(z, Z)$ with $Y = Y_0$. Explicitly,

$$D(\zeta) = \log \frac{\sqrt{\zeta^2 - 2\rho\zeta + 1} + \zeta - \rho}{1 - \rho}. \quad (79)$$

The analysis in Appendix B.2 shows that the probability distribution for x is Gaussian in this minimum distance, at least to leading order. Specifically, it is shown there that to within $O(\lambda^2)$,

$$\begin{aligned} P_X(s, x, y) = & \frac{1}{\sqrt{2\pi\lambda}} \frac{1}{yb(X) I^{3/2}} \exp\left\{-\frac{D^2}{2\lambda}\right\} \left\{1 + \frac{yb'(x) D}{2\sqrt{1-\rho^2} I}\right. \\ & + \lambda \left[-\frac{1}{8} + \frac{7}{16} \frac{1 - D \coth D}{D} \frac{yb'(x)}{\sqrt{1-\rho^2} I} \right. \\ & + \frac{3}{8} \frac{1 - \rho^2 \sinh D}{I D} \left(1 + \frac{yb'(x) D}{2\sqrt{1-\rho^2} I}\right) \\ & \left. \left. + \frac{1}{4} \frac{yb'(x)(\zeta - \rho)}{\sqrt{1-\rho^2} I} \right] + O(\lambda^2) \right\}, \end{aligned} \quad (80)$$

where

$$\begin{aligned} I(\zeta) &= \sqrt{\zeta^2 - 2\rho\zeta + 1} \\ &= \cosh D(\zeta) - \rho \sinh D(\zeta). \end{aligned} \quad (81)$$

Let us rewrite this expression in terms of the original variables:

$$\begin{aligned} P_F(\tau, f, \sigma) = & \frac{1}{\sqrt{2\pi\tau}} \frac{1}{\sigma b(F) I^{3/2}} \exp\left\{-\frac{D^2}{2\tau v^2}\right\} \left\{1 + \frac{\sigma b'(f) D}{2v\sqrt{1-\rho^2} I}\right. \\ & + \tau v^2 \left[-\frac{1}{8} + \frac{7}{16} \frac{1 - D \coth D}{D} \frac{\sigma b'(f)}{v\sqrt{1-\rho^2} I} \right. \\ & \left. \left. + \frac{3}{8} \frac{1 - \rho^2 \sinh D}{I D} \left(1 + \frac{\sigma b'(f) D}{2v\sqrt{1-\rho^2} I}\right) + \frac{\sigma b'(f)(\zeta - \rho)}{4v\sqrt{1-\rho^2} I} \right] \right\}. \end{aligned} \quad (82)$$

This is the desired asymptotic form of the marginal probability distribution.

5.2 Implied volatility

As the final step of our analysis, we shall derive, in the spirit of [10] explicit formulas for the equivalent normal volatility σ_n and the equivalent lognormal volatility

σ_{\ln} . To this order, we compare (82) with the normal distribution function:

$$n_F(\tau, f, \sigma_n) = \frac{1}{\sqrt{2\pi\tau}\sigma_n} \exp\left\{-\frac{(x-X)^2}{2\tau\sigma_n^2}\right\}. \quad (83)$$

The analysis is straightforward but somewhat tedious, and we defer it to Appendix B.3. The result of our calculations is the following asymptotic formula for σ_n in terms of the SABR parameters:

Likewise, comparing (82) with the lognormal distribution function

$$\sqrt{\frac{x}{2\pi\lambda X^3 y_t^2}} \exp\left(-\frac{(\log(x/X))^2}{2\lambda y_t^2} - \frac{\lambda y_t^2}{8}\right),$$

we obtain the asymptotic relationship:

5.3 Impact of boundary conditions at zero forward

5.4 Explicit models

Let us consider the special case of a stochastic CEV model, $b(f) = f^\beta$, with $0 < \beta \leq 1$. If $\beta = 1$, then

$$\zeta = \frac{v}{\sigma} \log \frac{f}{F}. \quad (84)$$

For $0 < \beta < 1$,

$$\zeta = \frac{v}{\sigma} \frac{f^\beta - F^\beta}{(1-\beta)}. \quad (85)$$

Consequently,

Another popular model is shifted lognormal model, $b(f) = f + a$, where $a > 0$. Then

$$\zeta = \frac{v}{\sigma} \log \frac{f+a}{F+a}, \quad (86)$$

and thus

A Heat equation on the Poincare plane

In this appendix we present an elementary derivation of the explicit representation of the Green's function for the heat equation on \mathbb{H}^2 . This explicit formula has been known for a long time (see e.g. [14]), and we include its construction here in order to make our calculations self-contained.

A.1 Lower bound on the Laplace-Beltrami operator

We shall first establish a lower bound on the spectrum of the Laplace-Beltrami operator on the Poincare plane. Let $\mathcal{H} = L^2(\mathbb{H}^2, d\mu_h)$ denote the Hilbert space of complex functions on \mathbb{H}^2 which are square integrable with respect to the measure (42). The inner product on this space is thus given by:

$$(\Phi|\Psi) = \int_{\mathbb{H}^2} \overline{\Phi(z)} \Psi(z) \frac{dx dy}{y^2}. \quad (87)$$

It is easy to verify that the Laplace-Beltrami operator Δ_h is self-adjoint with respect to this inner product.

Consider now the first order differential operator Q on \mathcal{H} defined by

$$Q = i \left(y \frac{\partial}{\partial y} - \frac{1}{2} \right) + y \frac{\partial}{\partial x}. \quad (88)$$

Its hermitian adjoint with respect to (87) is

$$Q^\dagger = i \left(y \frac{\partial}{\partial y} - \frac{1}{2} \right) - y \frac{\partial}{\partial x}, \quad (89)$$

and we verify readily that

$$\frac{1}{2} (QQ^\dagger + Q^\dagger Q) = -\Delta_h - \frac{1}{4}. \quad (90)$$

This implies that

$$\begin{aligned} (\Phi | -\Delta_h \Phi) &= \frac{1}{2} (\Phi | QQ^\dagger \Phi) + \frac{1}{2} (\Phi | Q^\dagger Q \Phi) + \frac{1}{4} (\Phi | \Phi) \\ &= \frac{1}{2} (Q^\dagger \Phi | Q^\dagger \Phi) + \frac{1}{2} (Q \Phi | Q \Phi) + \frac{1}{4} (\Phi | \Phi) \\ &\geq \frac{1}{4} (\Phi | \Phi), \end{aligned}$$

where we have used the fact that $(\Psi|\Psi) \geq 0$, for all functions $\Psi \in \mathcal{H}$. As a consequence, we have established that the spectrum of the operator $-\Delta_h$ is bounded from below by $\frac{1}{4}$! This fact was first proved in [14].

A.2 Construction of the Green's function

Let us now consider the the following initial value problem:

$$\begin{aligned} \frac{\partial}{\partial s} G_Z(s, z) &= \Delta_h G_Z(s, z), \\ G_Z(0, z) &= Y^2 \delta(z - Z), \end{aligned} \quad (91)$$

where $z, Z \in \mathbb{H}^2$. In addition, we require that

$$G_Z(s, z) \rightarrow 0, \quad \text{as } d(z, Z) \rightarrow \infty. \quad (92)$$

Note that, up to the factor of Y^2 in front of the delta function and a trivial time rescaling, this is exactly the initial value problem (64):

$$G_Z(s, z) = Y^2 K_Z(2s/\varepsilon, z). \quad (93)$$

The Green's function $G_Z(s, z)$ is also referred to as the heat kernel³ on \mathbb{H}^2 . The reason for inserting the factor of Y^2 in front of $\delta(z - Z)$ is that the distribution $Y^2\delta(z - Z)$ is invariant under the action (39) of the Lie group $SL(2, \mathbb{R})$. In fact, we verify readily that

$$Y^2\delta(z - Z) = \frac{1}{\pi} \delta(\cosh d(z, Z) - 1).$$

Now, since the initial value problem (92) is invariant under $SL(2, \mathbb{R})$, its solution must be invariant and thus a function of $d(z, Z)$ only. Let $r = \cosh d(z, Z)$, and write $G_Z(s, z) = \varphi(s, r)$. Then the heat equation in (92) takes the form

$$\frac{\partial}{\partial s}\varphi(s, r) = (r^2 - 1) \frac{\partial^2}{\partial r^2}\varphi(s, r) + 2r \frac{\partial}{\partial r}\varphi(s, r). \quad (94)$$

We have established above that the operator $-\Delta_h$ is self-adjoint on the Hilbert space \mathcal{H} , and its spectrum is bounded from below by $\frac{1}{4}$. Therefore, we shall seek the solution as the Laplace transform

$$\varphi(s, r) = \int_{1/4}^{\infty} e^{-s\lambda} L(\lambda, r) d\lambda \quad (95)$$

which yields the following ordinary differential equation:

$$(1 - r^2) \frac{d^2}{dr^2} L(\lambda, r) - 2r \frac{d}{dr} L(\lambda, r) + \lambda L(\lambda, r) = 0. \quad (96)$$

We write

$$\lambda = \nu(\nu + 1),$$

where

$$\begin{aligned} \nu &= -\frac{1}{2} \pm i\sqrt{\lambda - \frac{1}{4}} \\ &= -\frac{1}{2} \pm i\omega, \end{aligned}$$

³It is the integral kernel of the semigroup of operators generated by the heat equation.

and recognize in (96) the Legendre equation. Note that, as a consequence of the inequality $\lambda \geq \frac{1}{4}$, ω is real and $\operatorname{Re} \nu = -\frac{1}{2}$.

In the remainder of this appendix, we will use the well known properties of the solutions to the Legendre equation, and follow Chapters 7 and 8 of Lebedev's book on special functions [13]. The general solution to (96) is a linear combination of the Legendre functions of the first and second kinds, $P_{-1/2+i\omega}(r)$ and $Q_{-1/2+i\omega}(r)$, respectively:

$$L\left(\frac{1}{4} + \omega^2, r\right) = A_\omega P_{-1/2+i\omega}(r) + B_\omega Q_{-1/2+i\omega}(r). \quad (97)$$

As $d \rightarrow 0$ (which is equivalent to $r \rightarrow 1$),

$$Q_{-1/2+i\omega}(\cosh d) \sim \text{const} \log d, \quad (98)$$

which would imply that $\varphi(s, \cosh d)$ is singular at $d = 0$, for all values of $s > 0$. Since this is impossible, we conclude that $B_\omega = 0$. Note that, on the other hand,

$$P_{-1/2+i\omega}(1) = 1, \quad (99)$$

i.e. $P_{-1/2+i\omega}(\cosh d)$ is non-singular at $d = 0$.

We will now invoke the Mehler-Fock transformation of a function⁴:

$$\tilde{f}(\omega) = \int_1^\infty f(r) P_{-1/2+i\omega}(r) dr, \quad (100)$$

$$f(r) = \int_0^\infty \tilde{f}(\omega) P_{-1/2+i\omega}(r) \omega \tanh(\pi\omega) d\omega. \quad (101)$$

In particular, (99) implies that the Mehler-Fock transform of $\delta(r-1)$ is 1, and thus (remember that we need to divide $\delta(r-1)$ by π):

$$A_\omega = \frac{1}{2\pi} \tanh(\pi\omega).$$

Now, the Legendre function of the first kind $P_{-1/2+i\omega}(r)$ has the following integral representation:

$$P_{-1/2+i\omega}(\cosh d) = \frac{\sqrt{2}}{\pi} \coth(\pi\omega) \int_d^\infty \frac{\sin(\omega u)}{\sqrt{\cosh u - \cosh d}} du, \quad (102)$$

which is valid for all real ω . Therefore

$$L\left(\frac{1}{4} + \omega^2, \cosh d\right) = \frac{1}{\sqrt{2}\pi^2} \int_d^\infty \frac{\sin(\omega u)}{\sqrt{\cosh u - \cosh d}} du,$$

⁴Strictly speaking, we will deal with distributions rather than functions. A rigor oriented reader can easily recast the following calculations into respectable mathematics.

and we can easily carry out the integration in (95) to obtain

$$G_Z(s, z) = \frac{e^{-s/4}\sqrt{2}}{(4\pi s)^{3/2}} \int_{d(z, Z)}^{\infty} \frac{ue^{-u^2/4s}}{\sqrt{\cosh u - \cosh d(z, Z)}} du. \quad (103)$$

This is McKean's closed form representation of the Green's function of the heat equation on the Poincare plane [14].

Going back to the original normalization conventions of (64) yields formula (65).

B Some asymptotic expansions

In this appendix we collect a number of asymptotic expansions used in this paper.

B.1 Asymptotics of the McKean kernel

We shall establish a short time asymptotic expansion of the McKean kernel $G_Z(s, z)$ which plays a key role in the analysis of the Green's function of the SABR model.

In the right hand side of (103) we substitute $u = \sqrt{4sw + d^2}$:

$$G_Z(s, z) = \frac{e^{-s/4}\sqrt{2}}{4\pi^{3/2}\sqrt{s}} e^{-d^2/4s} \int_0^{\infty} \frac{e^{-w} dw}{\sqrt{\cosh \sqrt{4sw + d^2} - \cosh d}}.$$

Expanding the integrand in powers of s yields

$$\frac{1}{\sqrt{\cosh \sqrt{4sw + d^2} - \cosh d}} = \sqrt{\frac{d}{\sinh d}} \times \left(\frac{1}{\sqrt{2sw}} - \frac{d \coth d - 1}{4d^2} \sqrt{2sw} + O(s^{3/2}) \right).$$

Integrating term by term over w we find that

$$G_Z(s, z) = \frac{e^{-s/4}}{4\pi s} \exp\left(-\frac{d^2}{4s}\right) \times \sqrt{\frac{d}{\sinh d}} \left(1 - \frac{1}{4} \frac{d \coth d - 1}{d^2} s + O(s^2) \right),$$

and we thus obtain the following asymptotic expansion of the McKean kernel:

$$G_Z(s, z) = \frac{1}{4\pi s} \exp\left(-\frac{d^2}{4s}\right) \times \sqrt{\frac{d}{\sinh d}} \left(1 - \frac{1}{4} \left(\frac{d \coth d - 1}{d^2} + 1 \right) s + O(s^2) \right), \quad (104)$$

Taking the derivative of $G_Z(s, z)$ with respect of $d(z, Z)$ in the expansion above, we find that

$$\begin{aligned} \frac{\partial}{\partial d} G_Z(s, z) &= \frac{1}{4\pi s} \exp\left(-\frac{d^2}{4s}\right) \times \\ &\quad \sqrt{\frac{d}{\sinh d}} \left(-\frac{d}{2s} + \frac{1 - d \coth d}{2d} + O(s)\right). \end{aligned} \quad (105)$$

B.2 Laplace's method

Next we review the Laplace method (see e.g. [3], [2]) which allows one to evaluate approximately integrals of the form:

$$\int_0^\infty f(u) e^{-\phi(u)/\epsilon} du. \quad (106)$$

We use this method in order to evaluate the marginal probability distribution for the Green's function.

In the integral (106), ϵ is a small parameter, and $f(u)$ and $\phi(u)$ are smooth functions on the interval $[0, \infty)$ ⁵. We also assume that $\phi(u)$ has a unique minimum u_0 inside the interval with $\phi''(u_0) > 0$. The idea is that, as $\epsilon \rightarrow 0$, the value of the integral is dominated by the quadratic approximation to $\phi(u)$ around u_0 .

More precisely, we have the following asymptotic expansion. As $\epsilon \rightarrow 0$,

$$\begin{aligned} \int_0^\infty f(u) e^{-\phi(u)/\epsilon} du &= \sqrt{\frac{2\pi\epsilon}{\phi''(u_0)}} e^{-\phi(u_0)/\epsilon} \times \\ &\quad \left\{ f(u_0) + \epsilon \left[\frac{f''(u_0)}{2\phi''(u_0)} - \frac{\phi^{(4)}(u_0) f(u_0)}{8\phi''(u_0)^2} \right. \right. \\ &\quad \left. \left. - \frac{f'(u_0) \phi^{(3)}(u_0)}{2\phi''(u_0)^2} + \frac{5\phi^{(3)}(u_0)^2 f(u_0)}{24\phi''(u_0)^3} \right] + O(\epsilon^2) \right\}. \end{aligned} \quad (107)$$

To generate this expansion, we first expand $f(u)$ and $\phi(u)$ in Taylor series around u_0 to orders 2 and 4, respectively (keep in mind that the first order term in the expansion of $\phi(u)$ is zero). Then, expanding the regular terms in the exponential, we organize the integrand as $e^{-\phi''(u_0)(u-u_0)^2/2\epsilon}$ times a polynomial in ϵ . In the limit $\epsilon \rightarrow 0$, the integral reduces to calculating moments of the Gaussian measure; the result is (107). It is straightforward to compute terms of order higher than 1 in ϵ , even though the calculations become increasingly complex as the order increases.

⁵It can be an arbitrary interval.

Finally, let us state a slight generalization of (107), which we use below. In the integral (106), we replace $f(u)$ by $f(u) + \epsilon g(u)$. Then, as $\epsilon \rightarrow 0$,

$$\int_0^\infty [f(u) + \epsilon g(u)] e^{-\phi(u)/\epsilon} du = \sqrt{\frac{2\pi\epsilon}{\phi''(u_0)}} e^{-\phi(u_0)/\epsilon} \times \left\{ f(u_0) + \epsilon \left[g(u_0) + \frac{f''(u_0)}{2\phi''(u_0)} - \frac{\phi^{(4)}(u_0) f(u_0)}{8\phi''(u_0)^2} - \frac{f'(u_0) \phi^{(3)}(u_0)}{2\phi''(u_0)^2} + \frac{5\phi^{(3)}(u_0)^2 f(u_0)}{24\phi''(u_0)^3} \right] + O(\epsilon^2) \right\}. \quad (108)$$

This formula follows immediately from (107).

We shall now apply this formula in order to evaluate the integral (76). We find easily that the minimum Y_0 of the function

$$\phi(Y) = \frac{1}{2} \delta(z, Z)^2$$

is given by

$$Y_0 = y \sqrt{\zeta^2 - 2\rho\zeta + 1},$$

where

$$\zeta = \frac{1}{y} \int_X^x \frac{du}{b(u)}.$$

Also, we let $D(\zeta)$ denote the value of $\delta(z, Z)$ with $Y = Y_0$:

$$D(\zeta) = \log \frac{\sqrt{\zeta^2 - 2\rho\zeta + 1} + \zeta - \rho}{1 - \rho}.$$

and

$$I(\zeta) = \sqrt{\zeta^2 - 2\rho\zeta + 1}.$$

Let us now evaluate the various terms on the right hand side of (108). Note that the second derivative $\phi''(Y_0)$ of $\phi(Y)$ with respect to Y is

$$\phi''(Y_0) = \frac{D}{(1 - \rho^2) y^2 I \sinh D},$$

where we have suppressed the argument ζ in $D(\zeta)$ and $I(\zeta)$. Likewise,

$$\phi^{(3)}(Y_0) = -\frac{3D}{(1 - \rho^2) y^3 I^2 \sinh D},$$

and

$$\phi^{(4)}(Y_0) = \frac{3(1 - D \coth D)}{(1 - \rho^2)^2 y^4 I^2 \sinh^2 D} + \frac{12D}{(1 - \rho^2) y^4 I^3 \sinh D}.$$

Finally, in order to conform with the notation in (108), let us define

$$f(Y) = \sqrt{\frac{\delta}{\sinh \delta}} \left(1 - \frac{\delta}{\sinh \delta} q \right),$$

and

$$g(Y) = \sqrt{\frac{\delta}{\sinh \delta}} \left(\frac{1}{8} + \frac{\delta \coth \delta - 1}{8\delta^2} - \frac{\delta \coth \delta - 1}{2\delta \sinh \delta} q \right).$$

Then, after some manipulations we find that

$$f(Y_0) = \sqrt{\frac{D}{\sinh D}} \left(1 + \frac{yb'(x)D}{2\sqrt{1-\rho^2}I} \right),$$

$$f'(Y_0) = - \left(\frac{D}{\sinh D} \right)^{3/2} \frac{b'(x)(\zeta - \rho)}{2(1 - \rho^2)^{3/2} I^2},$$

$$\begin{aligned} f''(Y_0) &= \sqrt{\frac{D}{\sinh D}} \frac{1 - D \coth D}{2(1 - \rho^2) y^2 I D \sinh D} \left(1 + \frac{3yb'(x)D}{2\sqrt{1-\rho^2}I} \right) \\ &+ \left(\frac{D}{\sinh D} \right)^{3/2} \frac{b'(x)(\zeta - \rho)}{(1 - \rho^2)^{3/2} y I^3}, \end{aligned}$$

and

$$g(Y) = -\frac{1}{8} \sqrt{\frac{D}{\sinh D}} \left(1 + \frac{D \coth D - 1}{D^2} \left(1 + \frac{2yb'(x)D}{\sqrt{1-\rho^2}I} \right) \right).$$

Putting everything together we find that

$$\begin{aligned} P_X(s, x, y) &= \frac{1}{\sqrt{2\pi\lambda}} \frac{1}{yb(X) I^{3/2}} \exp \left\{ -\frac{D^2}{2\lambda} \right\} \left\{ 1 + \frac{yb'(x)D}{2\sqrt{1-\rho^2}I} \right. \\ &+ \lambda \left[-\frac{1}{8} + \frac{7}{16} \frac{1 - D \coth D}{D} \frac{yb'(x)}{\sqrt{1-\rho^2}I} \right. \\ &+ \frac{3}{8} \frac{1 - \rho^2 \sinh D}{I D} \left(1 + \frac{yb'(x)D}{2\sqrt{1-\rho^2}I} \right) \\ &\left. \left. + \frac{1}{4} \frac{yb'(x)(\zeta - \rho)}{\sqrt{1-\rho^2}I} \right] + O(\lambda^2) \right\}, \end{aligned}$$

as stated in Section 5.

B.3 Normalizing the SABR distribution

In this appendix, we shall relate the normal cumulative distribution function

$$\begin{aligned} \int_{\tilde{x}}^{\infty} n_X(s, x, y_n) df &= \frac{1}{\sqrt{2\pi y_n^2}} \int_{\tilde{x}}^{\infty} \exp\left\{-\frac{(x-X)^2}{2s y_n^2}\right\} dx \\ &= \frac{1}{2} \operatorname{erfc}\left(\frac{\tilde{x}-X}{\sqrt{2s y_n^2}}\right), \end{aligned} \quad (109)$$

where $\operatorname{erfc}(u)$ denotes the standard complementary error function,

$$\operatorname{erfc}(u) = \frac{2}{\sqrt{\pi}} \int_u^{\infty} e^{-x^2} dx,$$

to the cumulative distribution function of $P_X(s, x, y)$:

$$\begin{aligned} \int_{\tilde{x}}^{\infty} P_X(s, x, y) dx &= \frac{1}{\sqrt{2\pi\lambda} y b(X)} \int_{\tilde{x}}^{\infty} \frac{1}{I^{3/2}} \exp\left\{-\frac{D^2}{2\lambda}\right\} \times \\ &\quad \left\{ 1 + \frac{y b'(x) D}{2\sqrt{1-\rho^2} I} + \lambda \left[-\frac{1}{8} + \frac{7}{16} \frac{1-D \coth D}{D} \frac{y b'(x)}{\sqrt{1-\rho^2} I} \right. \right. \\ &\quad \left. \left. + \frac{3}{8} \frac{1-\rho^2 \sinh D}{I D} \left(1 + \frac{y b'(x) D}{2\sqrt{1-\rho^2} I} \right) + \frac{y b'(x) (\zeta - \rho)}{4\sqrt{1-\rho^2} I} \right] \right\} dx. \end{aligned}$$

We have temporarily placed a tilde over today's forward x , in order not to confuse it with the integration variable. We now substitute a new variable in the integral above,

$$h = h(x) = D \left(\frac{1}{y} \int_X^x \frac{du}{b(u)} \right),$$

and note that the Jacobian has a simple and helpful form:

$$\frac{dh}{dx} = \frac{1}{y b(x) I} > 0.$$

In particular, the function $h = h(x)$ is invertible, and we can write $x = x(h)$. This yields

$$\begin{aligned} \int_{\tilde{x}}^{\infty} P_X(s, x, y) dx &= \frac{1}{\sqrt{2\pi\lambda} y b(X)} \int_{h(\tilde{x})}^{\infty} \exp\left\{-\frac{h^2}{2\lambda}\right\} \frac{b(x(h))}{\sqrt{i(h)}} \times \\ &\quad \left\{ 1 + \frac{y b'(x(h)) h}{2\sqrt{1-\rho^2} i(h)} + \lambda \left[-\frac{1}{8} + \frac{7}{16} \frac{1-h \coth h}{h} \frac{y b'(x(h))}{\sqrt{1-\rho^2} i(h)} \right. \right. \\ &\quad \left. \left. + \frac{3}{8} \frac{1-\rho^2 \sinh h}{i(h) h} \left(1 + \frac{y b'(x(h)) h}{2\sqrt{1-\rho^2} i(h)} \right) + \frac{y b'(x(h)) j(h)}{4\sqrt{1-\rho^2} i(h)} \right] \right\} dh, \end{aligned}$$

where

$$\begin{aligned} i(h) &= I(\zeta) = \cosh h - \rho \sinh h, \\ j(h) &= \zeta - \rho = \sinh h - \rho \cosh h. \end{aligned}$$

Inspection of the integral on the right hand side of the equation above shows that in order to calculate it to within $O(\lambda^2)$, we need to Taylor expand the $O(1)$ terms in the integrand out to order 2 in h around 0, and we replace the $O(\lambda)$ terms by their value at $h = 0$. The result is the equality:

$$\begin{aligned} & \int_{\tilde{x}}^{\infty} P_X(s, x, y) dx \\ &= \frac{1}{\sqrt{2\pi\lambda}} \int_{h(\tilde{x})}^{\infty} \exp\left\{-\frac{h^2}{2\lambda}\right\} \left\{1 + \left(\frac{\rho}{2} + \frac{yb'(X)}{2\sqrt{1-\rho^2}} + b'(X)\right) h \right. \\ &+ \left(-\frac{2-3\rho^2}{4} + y^2b(X)b''(X) + yb'(X)(\rho + yb'(X)) \left(1 + \frac{1}{\sqrt{1-\rho^2}}\right)\right) \frac{h^2}{2} \\ &+ \left.\lambda \left[\frac{2-3\rho^2}{8} - \frac{\rho yb'(X)}{4\sqrt{1-\rho^2}}\right]\right\} dh + \dots \end{aligned}$$

We now suppress the tilde, and rewrite the last equation in the equivalent form:

References

- [1] Beardon, A. F.: *The Geometry of Discrete Groups*, Springer Verlag (1983).
- [2] Bender, C. M., and Orszag, S. A.: *Advanced Mathematical Methods for Scientists and Engineers*, Springer Verlag (1999).
- [3] Bleistein, N., and Handelsman, R. A.: *Asymptotic Expansions of Integrals*, Dover Publications (1986).
- [4] Duffie, D., Pan, J., and Singleton, K.: Transform analysis and asset pricing for affine jump diffusions, *Econometrica*, **68**, 1343 - 1377 (2000).
- [5] Elworthy, K. D.: *Geometric Aspects of Diffusions on Manifolds*, Ecole d'Ete de Probabilites de Saint Flour XVII, July 1987.
- [6] Emery, M.: *Stochastic Calculus in Manifolds*, Springer Verlag (1989).
- [7] Guenther, R. B., and Lee, J. W.: *Partial Differential Equations of Mathematical Physics and Integral Equations*, Prentice Hall (1988).

- [8] Hagan, P. S., Kumar, D., Lesniewski, A., and Woodward, D. E: Managing smile risk, *Wilmott Magazine* (2003).
- [9] Hagan, P. S., and Lesniewski, A.: in preparation.
- [10] Hagan, P. S., and Woodward, D. E: Equivalent Black Volatilities, *App. Math. Finance*, **6**, 147-157 (1999).
- [11] Heston, S.: A closed form solution for options with stochastic volatility with applications to bond and currency options, *Review of Financial Studies*, **6**, 327-343 (1993).
- [12] Hsu, E. P.: *Stochastic Analysis on Manifolds*, American Mathematical Society (2002).
- [13] Lebedev, N. N.: *Special Functions and their Applications*, Dover Publications (1972)
- [14] McKean, H. P.: An upper bound to the spectrum of Δ on a manifold of negative curvature, *J. Diff. Geom.*, **4**, 359-366 (1970).
- [15] Lewis, A. L.: *Option Valuation under Stochastic Volatility*, Finance Press (2000).
- [16] Molchanov, S. A.: Diffusion processes and Riemannian geometry, *Russian Math. Surveys*, **30** (1975) 1 - 63.
- [17] Varadhan, S. R. S: On the behavior of the fundamental solution of the heat equation with variable coefficients, *Comm. Pure Appl. Math.* **20** (1967) 431 - 455.
- [18] Varadhan, S. R. S: Diffusion processes in a small time interval, *Comm. Pure Appl. Math.* **20** (1967) 659 - 685.